

The relativistic glider

Eduardo Guéron*

Centro de Matemática, Computação e Cognição,
Universidade Federal do ABC, 09210-170, Santo André, SP, Brazil

Ricardo A. Mosna†

Instituto de Matemática, Estatística e Computação Científica,
Universidade Estadual de Campinas, C.P. 6065, 13083-859, Campinas, SP, Brazil.

We present a purely relativistic effect according to which *asymmetric* oscillations of a quasi-rigid body slow down or accelerate its fall in a gravitational background.

PACS numbers: 04.20.-q, 04.25.-g, 45.40.-f

The motion of extended bodies in General Relativity has been a source of unexpected phenomena up to the present. Although the problem in its full generality is notoriously complicated [1], approximated analysis of quasi-rigid bodies already discloses interesting effects. A remarkable example and quite surprising result was recently presented by Wisdom, who demonstrated that cyclic changes in the shape of a quasi-rigid body may lead to net translation in curved spacetimes [2]. Since this takes place in the absence of external forces, one can portray the body as swimming in spacetime. Later, this effect was shown to dominate over the so-called “swinging” (Newtonian) effect, whose formulae are well known, for high enough oscillation frequencies [3]. The latter is a direct expression of the fact that expanding and contracting a body in a nonuniform gravitational field gives rise to a net work on it (due to tidal forces). However, this effect is increasingly suppressed for higher frequencies, since the gravitational potential felt by the body is then nearly the same through its complete cycle (see Eq. 5 of [3]).

Here we present a purely relativistic effect which (like the swimming and unlike the swinging effect) is not suppressed for high frequencies, but (like the swinging and unlike the swimming effect) is not related to a gravitational geometric phase. It comes about by letting a quasi-rigid body perform *asymmetric* oscillations in a gravita-

tional background. We show that, as a result, the body slows down or accelerates its fall depending on how asymmetric its internal oscillations are.

We propose a simple model to investigate this effect: two test particles of mass m connected by a massless strut whose length varies periodically, i.e., an oscillating dumbbell. The strut is mathematically implemented as a constraint between the particles. Physically, and following Wisdom’s idea (see last section of [2]), the active strut should be constantly and locally monitored in order to guarantee that the constraints are maintained.

A test particle under the influence of a spherically symmetric gravitational field moves, in the context of General Relativity, in Schwarzschild spacetime. Its equations of motion follow from the Lagrangian

$$\mathcal{L}_p = -mc \sqrt{\left(1 - \frac{2GM}{c^2 r_p}\right) c^2 \left(\frac{dt}{ds}\right)^2 - \frac{1}{1 - \frac{2GM}{c^2 r_p}} \left(\frac{dr_p}{ds}\right)^2}, \quad (1)$$

where m is the mass of the test particle and s is an evolution parameter (we assume that the body falls vertically towards the gravitational source). For the dumbbell, we choose the coordinate time t as the evolution parameter and take the relevant Lagrangian as

$$\mathcal{L}_d = -mc \sqrt{\left(1 - \frac{2GM}{c^2 r_1}\right) c^2 - \frac{1}{1 - \frac{2GM}{c^2 r_1}} \left(\frac{dr_1}{dt}\right)^2} - mc \sqrt{\left(1 - \frac{2GM}{c^2 r_2}\right) c^2 - \frac{1}{1 - \frac{2GM}{c^2 r_2}} \left(\frac{dr_2}{dt}\right)^2} \Bigg|_{r_2(t)=r_1(t)+l(t)}, \quad (2)$$

where $l(t)$ is the constraint as seen by the observer at infinity. In the Newtonian limit, $l(t)$ represents the length of the strut connecting the particles at time t . This implementation of the constraint clearly embodies considerable simplification (for instance, it fails to be covariant) [4]. However, it is important to note that, by choosing a pe-

riodic $l(t)$ such that $l(0) = l(T) = 0$, all observers will agree that the worldlines of both particles meet at the end of each period. In the following, we numerically solve the equations of motion coming from (2) for a family of constraint functions $l_\alpha(t)$ whose members always meet this condition. Of course, different observers will disagree on

the symmetry and shape of a given function $l(t)$ as measured by them.

With these considerations in mind, we compute the fall of the oscillating dumbbell and compare it with that of the single particle (the particular case when $l(t) \equiv 0$), both with the same initial conditions. Given that the cyclic changes considered here affect a single degree of freedom, we can anticipate that the swimming effect will not arise in this case [5]. Therefore the sought after relativistic effect will be isolated once we work in a regime for which (Newtonian) tidal effects are negligible.

We numerically solved the Euler-Lagrange equations associated with (2) for the family $\{l_\alpha(t)\}$ portrayed in Fig. 1a. The asymmetry parameter α , ranging from -1 to 1 , quantifies how much $l_\alpha(t)$ fails to be symmetric about its peak: the function corresponding to $\alpha = 0$ has its peak at $t = T/2$, while a generic $l_\alpha(t)$ attains its maximum at $t = \frac{1+\alpha}{2} T$ [6]. For each value of α , the equations of motion of the oscillating dumbbell and of the point mass were integrated, with $r_p(0) = r_1(0) \equiv R$ and $\dot{r}_p(0) = \dot{r}_1(0) = 0$. Their separation $\delta r = r_1(T) - r_p(T)$ after one period then yields the net displacement of the dumbbell. Representative results, for $R = 120$ and $\delta l = 5 \times 10^{-3}$, in units of GM/c^2 , are shown in Fig. 1b, where $\delta r \times 10^9$ was plotted against the dumbbell oscillation frequency $\omega = 1/T$. The relativistic case, corresponding to (1) and (2), is represented by solid lines, while its Newtonian counterpart is depicted by dashed lines.

Fig. 1b shows that the Newtonian δr is always negative and that it becomes negligible when the frequency is large enough (as expected from our earlier discussion). It is important to note that this happens *irrespective of the value of the asymmetry parameter α* . This is in marked contrast to the relativistic case. Indeed, while the relativistic curve for the symmetric case ($\alpha = 0$) is surrounded by Newtonian curves, those corresponding to positive values of α lead to a *positive* net displacement, a purely relativistic effect. Therefore, a relativistic robot who cannot swim may still soften its fall by repeatedly stretching and shrinking its body in an asymmetric fashion. Note that the relativistic δr does not become negligible for large values of ω , and that its sign and magnitude in this regime are determined by α .

It is worth noting that the beginning of the plateaus in Fig. 1b occurs in a frequency domain which is fairly independent on the oscillation amplitude δl [7]. By keeping δl small one can thus obtain the effect in hand for reasonably small velocities. It is interesting to note that, by Taylor expanding the Lagrangian (2) in $\dot{r}_1$ and $\dot{r}_2$ while keeping the remaining (position-dependent) terms unperturbed, one needs to retain terms at least up to fourth order in v/c to capture the effect under consideration. This should be compared to Wisdom's swimming effect, which already takes place when the relevant Lagrangian is expanded up to second order in v/c . The effect presented here thus arises from an intricate coupling of the gravitational field to the velocities of the particles.




FIG. 1: (a) Family of constraint functions $l_\alpha(t)$ [6]. (b) Net displacement $\delta r = r_1(T) - r_p(T)$ of the dumbbell, relative to the falling point mass, after one period. The initial conditions are set by $\dot{r}_p(0) = \dot{r}_1(0) = 0$, $r_p(0) = r_1(0) = 120$ and $\delta l = 5 \times 10^{-3}$ (in units of GM/c^2). The relativistic (Newtonian) case is represented by solid (dashed) lines. The plot shows that the Newtonian δr is always negative and that it is negligible for $\omega \gtrsim 0.03$ c^3/GM , irrespective of α . In contrast, we see that the relativistic δr crucially depends on the asymmetry parameter α , and does attain positive values (for positive α). Causality problems (due to large velocities) only start to appear at much higher frequencies, not shown above.

From some numerical analysis, we estimate that the net displacement after one period, in the large frequency regime (corresponding to the plateaus in Fig. 1), is given by

$$\delta r \approx \Gamma_\alpha \frac{\delta l^2}{R^2} \frac{GM}{c^2}, \quad (3)$$

where δl is the amplitude of $l_\alpha(t)$ and Γ_α is a dimensionless factor associated with the asymmetry parameter α , see Fig. 2. This expression was obtained from numerical simulation employing a large range of values for R and δl , going from 50 to 10^7 and $10^{-7}R$ to $10^{-2}R$, respectively. Equation 3 shows that the effect under consideration can be orders of magnitude higher than the swimming effect, which is proportional to $\delta l^2/R^3$ [8]. This result is somewhat surprising since, as discussed above, the swimming effect is already present when the Lagrangian is expanded up to second order in the velocities, while the effect presented here needs at least a fourth-order expansion in v/c .




FIG. 2: Adimensional scale factor Γ_α as a function of the asymmetry parameter α .

It is instructive to calculate the net displacement δr in terms of the coordinate distance D traveled by the falling point mass during the period T . Using the rough estimate $D \sim gT^2$, with $g = GM/R^2$, we immediately see that

$$\delta r/D \sim (\delta l \omega / c)^2 \quad (4)$$

in this regime. In this way, if the dumbbell is engineered so that $\delta l \omega \sim 0.01c$, we get $\delta r/D \sim 10^{-4}$. Therefore, such result may be within reach of experimental verification. This would open the door to new tests of General Relativity by means of local experiments [9].

We close by noting that, if $\delta r/D$ were to reach 1, the dumbbell would be able to float in space. However, this is apparently not possible, since numerical simulations indicate that the value attained by $\delta r/D$ is typically limited to 15%. Therefore, it seems that, although this mechanism cannot be employed to fly, it might be conceivably used as a relativistic paraglider to slightly soften the landing on an atmosphereless planet.

Acknowledgments

The authors thank G. E. A. Matsas, S. R. Oliveira and Alberto Saa for fruitful discussions. EG is grateful to P. S. Letelier for the technical support. RAM acknowledges financial support from FAPESP.

-
- [1] W. G. Dixon, Proc. R. Soc. Lond. A **314**, 499 (1970);
W. G. Dixon, Nuovo Cimento **34**, 317 (1964).
 - [2] J. Wisdom, Science **299**, 1865 (2003).
 - [3] E. Guéron, C. A. S. Maia and G. E. A. Matsas, Phys.Rev. D**73**, 024020 (2006).
 - [4] A related (and also simplified) quasi-rigidity condition could be defined by imposing that the geodesic length of constant t slices connecting the worldlines of the particles be given by the constraint function $l(t)$. The obtained results are essentially the same for either of these choices.
 - [5] It is well-known that any geometric phase effect (including the swimming effect, Berry's phase, Hannay angle, etc) can only happen if at least two parameters are varied along a (nontrivial) loop in the appropriate parameter space. See, e.g., the book "A. Shapere and F. Wilczek, *Geometric Phases in Physics*, World Scientific, Singapore (1989)".
 - [6] The functional form of the constraint functions in Fig. 1a

is given by $l_\alpha(t) = \delta l \exp \left[\frac{(1+\alpha-2\omega t)^2}{(1+\alpha^2)\omega t(-1+\omega t)} \right]$, where $\omega = 1/T$ and the overall normalization was chosen so that the amplitude of $l_\alpha(t)$ is δl . Other families $\{l_\alpha(t)\}$ with diverse shapes were also tested, with no appreciable changes in the results.

- [7] Numerical simulations indicate that a variation in δl of five orders of magnitude leads to a change of less than 1% in the characteristic frequency for which the corresponding plateau begins.
- [8] See eq. (20) of reference [2].
- [9] We note that, if M is the Earth's mass, then $R \sim 10^9 GM/c^2$ at Earth's surface. Although we could numerically confirm Eq. (3) only for $50 GM/c^2 \lesssim R \lesssim 10^7 GM/c^2$ (due to limitations of our integration routine), there is no indication that it would break down at such scale.